


# Vitreomacular Adhesion

Vitreomacular adhesion, also known as vitreomacular traction syndrome, refers to an attachment between the vitreous gel which fills the eyeball and the central part of the retina which lines the back of the eye. If the vitreous gel separates from the retina except for a persistent central attachment, then the gel can physically pull on the retina and cause swelling or deformation which results in blurred or distorted vision.

## How does vitreomacular adhesion occur?

At birth, the vitreous gel is clear and firm. The surface of the gel is attached to the retina, as shown in the first image below. With age, the gel begins to liquefy and soften. Eventually, in most eyes, the surface of the gel separates cleanly from the retina, creating a posterior vitreous detachment, as shown in the second image below. However, in some eyes the attachment between the gel and the retina is strong enough that the gel separates only partially, leaving a small area of attachment as shown in the third image below. This attachment can pull on the retina cause various deformations.



## How is vitreomacular adhesion diagnosed?

Your retinal surgeon may order diagnostic tests in the office to identify and characterize the adhesion between the vitreous gel and the retina.

*Optical coherence tomography* (OCT) is a retinal scan that measures abnormalities in very high resolution. In some cases of vitreomacular adhesion, *fluorescein angiography* imaging is helpful in identifying blood vessel leakage associated with the traction exerted by the vitreous gel.

**BayAreaRetinaAssociates**

*Diseases and Surgery of the Retina and Vitreous*



800-5-RETINA (800-573-8462)

<http://www.bayarearetina.com>

Allen Verne MD | Craig Leong MD | Stewart Daniels MD | Subhransu Ray MD, PhD | Daniel Ting MD, PhD | Tushar Ranchod MD | Roger Goldberg MD, MBA

ANTIOCH

CASTRO VALLEY

FREMONT

OAKLAND

PLEASANTON

SAN LEANDRO

VALLEJO

WALNUT CREEK

## What treatments are available for vitreomacular adhesion?

**Observation.** In some cases, vitreomacular adhesions will resolve on their own and treatment can be avoided. Spontaneous resolution is difficult to predict, and so observation for at least a short period of time is usually appropriate before initiating any type of treatment.

**Intravitreal ocriplasmin injection.** Jetra (ocriplasmin) was FDA approved in 2012 for treatment of vitreomacular adhesion. This clear liquid injection is given in the office after the eye is numbed. The odds of resolution with this treatment vary depending on the exact characteristics of the adhesion. If the adhesion does not resolve with ocriplasmin treatment, then vitrectomy surgery is usually required to treat the persistent vitreomacular adhesion.



**Micro-incisional / sutureless vitrectomy surgery.** Micro-incisional vitrectomy surgery is performed in the operating room as a same-day procedure. The surgery is able to successfully separate the vitreomacular adhesion more than 90% of the time. In some cases, a gas bubble is placed in the eye and face-down position is required in order to achieve the best visual outcome after surgery.

## Will my vision improve after treatment?

Visual improvement depends on several factors. Without physical separation of the adhesion, vision is unlikely to improve. If the adhesion separates (spontaneously or with treatment), the vision is likely but not guaranteed to improve. Visual improvement is less likely if the retinal deformation is very severe or has been present for a long period of time.

If the adhesion separates but causes a full thickness macular hole, then vision may actually worsen, and treatment is then required to close the macular hole.

► Optical coherence tomography images showing a vitreomacular adhesion (left) and an adhesion resulting in macular hole (right).



**BayAreaRetinaAssociates**

*Diseases and Surgery of the Retina and Vitreous*



800-5-RETINA (800-573-8462)

<http://www.bayarearetina.com>

Allen Verne MD | Craig Leong MD | Stewart Daniels MD | Subhransu Ray MD, PhD | Daniel Ting MD, PhD | Tushar Ranchod MD | Roger Goldberg MD, MBA

ANTIOCH    CASTRO VALLEY    FREMONT    OAKLAND    PLEASANTON    SAN LEANDRO    VALLEJO    WALNUT CREEK

Bay Area Retina Associates is a group practice of retinal surgeons. All members of the group are board certified by the American Academy of Ophthalmology and have completed fellowship training in vitreoretinal surgery. BARA surgeons have expertise in the treatment of retinal detachment, diabetic retinopathy, age-related macular degeneration, macular hole, epiretinal membrane, and retinal vascular disease. BARA physicians see patients in eight offices and perform surgery at several hospitals and surgery centers around the East Bay.